

Correlation functions of the global E_8 symmetry currents in the Heterotic 5-brane theory.

Morten Krogh

*Department of Physics, Jadwin Hall, Princeton University,
Princeton, NJ 08544, USA*

We consider the 5-brane placed at one end of the world in the Heterotic $E_8 \times E_8$ theory. The low energy theory is a 6 dimensional $(1, 0)$ superconformal theory with E_8 as a global symmetry. We calculate the two-point correlator of the E_8 current in 6 dimensions and in 4 dimensions after compactification on $\mathbf{T}^2$. This correlator is derived in 3 different ways: From field theory, from 11 dimensional supergravity and from F-theory.

This paper is a written version of the talk given at the 1998 Trieste Conference on Superfivebranes and Physics in 5+1 Dimensions. The talk presented the work of ¹, where a more elaborate discussion can be found.

1 Introduction

In the past two years, many examples of nontrivial IR fixed points in various dimensions have emerged. Some of the most exciting ones are the 5+1D chiral theories. The first of such theories with $\mathcal{N} = (2, 0)$ SUSY has been discovered in ² as a sector of type-IIB compactified on an A_1 singularity. A dual realization was found in ³ as the low-energy description of two 5-branes of M-theory. Another theory of this kind arises as an M-theory 5-brane approaches the 9-brane ^{8,10}. When the distance between the 5-brane and 9-brane is zero, the low-energy is described by a nontrivial 5+1D fixed point. This theory is chiral with $\mathcal{N} = (1, 0)$ SUSY and a global E_8 symmetry. In ^{10,5} more examples of $\mathcal{N} = (1, 0)$ theories have been given. We will use the terminology of ⁵ and call the E_8 theory V_1 . Many other 5+1D theories have been recently constructed in ^{14,15}.

String theory is a powerful tool to study such theories. The idea is to identify a dual description such that quantum corrections of the original theory appear at the classical level of the dual ¹⁹. The toroidal compactification of the $\mathcal{N} = (1, 0)$ 6D theory (and hence 4D $\mathcal{N} = 2$ QCD) can be studied using the brane-probe technique discovered in ^{13,12}. The world-volume theory on a brane probe in a heterotic string vacuum (which is quantum mechanically corrected) is mapped by duality to a world-volume theory on a brane inside a curved background which is not quantum mechanically corrected. This allows one to

determine the low-energy behavior in 4D. At the origin of the moduli space one obtains an IR fixed point with E_8 global symmetry.

The purpose of the present work is to extract information about the local operators of such theories. The E_8 theory V_1 has a local E_8 current $j_\mu^a(x)$ ($a = 1 \dots 248$ and $\mu = 0 \dots 5$). We will be interested in the correlator $\langle j_\mu^a(x) j_\nu^b(0) \rangle$. The strategy will be to couple the theory to a weakly coupled E_8 gauge theory and calculate the effect of V_1 on the E_8 coupling constant. We will study the question both for the 5+1D theory and for the 3+1D conformal theories. We will present three methods for evaluating the correlator. The first method is purely field-theoretic and applies to the 3+1D theories. Deforming the theory with a relevant operator one can flow to the IR where a field-theoretic description of $SU(2)$ or $U(1)$ with several quarks⁹ can be found. This will allow us to determine the correlator as a function on the moduli space. From this function we can deduce the high-energy behavior of the correlator and find out how many copies of the E_8 theory can be gauged with an E_8 SYM before breaking asymptotic freedom.

The other two methods for determining the correlators involve M-theory and F-theory. The gravitational field of a 5-brane of M-theory which is close to a 9-brane changes the local metric on the 9-brane. After compactification on a large $K3$ this implies that the volume of the $K3$ at the position of the 9-brane is affected by the distance from the 5-brane (see⁴). This can be interpreted as a dependence of the E_8 coupling constant on the VEV which specifies the position of the 5-brane. From this fact we can extract the current correlator. The third method involves the F-theory¹¹ realization of the E_8 theory^{10,16,6}. The V_1 theory is obtained in F-theory compactifications on a 3-fold by blowing up a point in the (two complex dimensional) base. By studying the effect of the size of the blow-up on the size of the 7-brane locus we can again determine the dependence of the E_8 coupling constant on the VEV.

The paper is organized as follows. In section (2) we calculate the current-current correlators in 3+1D using field theory arguments and we argue that 10 copies of the E_8 theory can be coupled to a gauge field. In section (3) we study the effect of a 5-brane on the volume of a 9-brane in M-theory and deduce the correlator from this setting. In section (4) we present the F-theory derivation. In section (5) we conclude.

2 Field theory derivation of the 4 dimensional correlator

In this section we will derive the form of the E_8 current-current correlator for the E_8 conformal theory and as a result we will argue that in 4D one can couple up to 10 copies of the E_8 theory to a $\mathcal{N} = 2$ E_8 Yang-Mills gauge theory.

We start with the E_8 conformal theory in 4 dimensions whose Seiberg-Witten curve is given by⁹:

$$y^2 = x^3 + u^5, \quad (1)$$

u parameterizing the moduli space of the Coulomb branch. We are looking for an expression of the form

$$\langle j_\mu^a(q) j_\nu^b(-q) \rangle = (q^2 \eta_{\mu\nu} - q_\mu q_\nu) \delta^{ab} f(q^2, u, \Lambda), \quad a, b = 1 \dots 248, \quad (2)$$

where q is the momentum and Λ is some fixed UV-cutoff. This UV-cutoff is not physical. It is just an artifact of the Fourier transform. The space-time correlator $\langle j_\mu^a(x) j_\nu^b(y) \rangle$ does not require a cutoff.

To determine the form of f in 2 for $q^2 = 0$ we can couple the E_8 SCFT to a weakly coupled E_8 gauge field and ask how the E_8 coupling constant changes as a function of u . When the E_8 coupling constant is very small the coupling does not change the curve 1 by much. For a generic value of u the massless modes of the E_8 SCFT are neutral under the global E_8 and the charged matter has a typical energy of order $u^{1/6}$. The $\langle jj \rangle$ correlator will modify the low energy E_8 coupling constant to the form

$$\frac{1}{g(u)^2} = \frac{1}{(g_0)^2} + f(q^2 = 0, u, \Lambda),$$

where g_0 is the bare coupling constant. On the other hand, standard renormalization arguments require that it should be possible to re-absorb the Λ dependence in the bare coupling constant. Thus, dimensional analysis restricts the form of $f(0, u, \Lambda)$ to

$$f(0, u, \Lambda) = c \log \left(\frac{\Lambda}{|u|^{1/6}} \right). \quad (3)$$

To determine c we deform the theory by adding a relevant operator to its (unknown) Lagrangian such as to break the global E_8 symmetry down to D_4 ($SO(8)$) by putting Wilson lines on the torus. The advantage is that the D_4 conformal fixed point can be analyzed in standard field-theory. It is the IR free theory of $SU(2)$ coupled to 4 massless quarks⁹.

The global E_8 of the original theory has been broken by the operators to a global $SO(8)$. For the $SO(8)$ theory we can ask what is

$$\langle j_\mu^A(q) j_\nu^B(-q) \rangle, \quad A, B = 1 \dots 28, \quad (4)$$

where A, B are $SO(8)$ indices. The point is now that we can calculate this correlator for the $SO(8)$ theory from field theory. The relevant field theory is

$SU(2)$ gauge theory with 4 quarks. From this correlator we can extract the original E_8 correlator. The details can be found in¹. We conclude that for the E_8 theory

$$\langle j_\mu^a(q) j_\nu^b(-q) \rangle = -\frac{3C(\text{fund.})}{4\pi^2} \delta^{ab} (q^2 \eta_{\mu\nu} - q_\mu q_\nu) \log \left| \frac{(\Lambda \lambda^2)^{1/3}}{u^{1/6}} \right|. \quad (5)$$

This means that the value of c in 3 is

$$c = -3 \frac{C(\text{fund. of SO}(8))}{4\pi^2} = -\frac{1}{10} \times \frac{C_2(E_8)}{4\pi^2}.$$

This value of c implies that 10 copies of the E_8 SCFT can be coupled to an E_8 SYM.¹

3 Derivation from M-theory

The system of the $(1,0)$ E_8 theory (V_1) coupled to E_8 SYM can be realized in M-theory as a 5-brane which is close to the 9-brane. The modes of the V_1 theory come from the 5-brane bulk and from membranes stretched between the 5-brane and 9-brane while the E_8 SYM comes from the 9-brane bulk. Let us compactify on $K3 \times \mathbf{T}^2$.

The effect that we are trying to study corresponds to the following question. The gravitational field of the 5-brane affects the metric at the position of the 9-brane. Thus, as we change the distance of the 5-brane from the 9-brane the volume of the $K3$ changes as a function of x ⁴. The volume of $K3 \times \mathbf{T}^2$ is related to the 3+1D E_8 coupling constant. In field-theory, this is interpreted as a running of the E_8 coupling constant as a result of the change of the VEV of the V_1 theory.

We apply the general setting and formulae of⁴ to the case where the distance of the 5-brane from the 9-brane is much smaller than the compactification scale of $K3 \times \mathbf{T}^2$ and calculate the effect.

We must also mention that the after compactification of the system of a 5-brane and 9-brane on $\mathbf{S}^1$ we get 4-branes near 8-branes. This setting has been studied in¹⁷, in the context of brane probes, where a related effect is observed. The position of the probe affects the value of a classical field, in that case the dilaton, which is then re-interpreted as a 1-loop effect in field theory. In fact, the relation between the classical supergravity calculation and the 1-loop field-theory calculation follows from perturbative string-theory. The 1-loop result is a loop of DD strings connecting the 4-brane to the 8-brane while the classical supergravity result is the same diagram viewed from the closed string channel.

3.1 Geometrical setup and review

In this section we will examine the theory of a 5-brane in M-theory on $\mathbb{R}^{5,1} \times K3 \times \mathbf{S}^1/\mathbb{Z}_2$ and review some relevant facts from^{7, 4}.

The geometric setup is as follows. The coordinates $(x^1, x^2, \dots, x^6)$ parameterize $\mathbb{R}^{5,1}$, (x^7, x^8, x^9, x^{10}) parameterize K3 and finally x^{11} parameterize $\mathbf{S}^1/\mathbb{Z}_2$. All 5-branes have their world-volume along $\mathbb{R}^{5,1}$ and are located at a point in $K3 \times \mathbf{S}^1/\mathbb{Z}_2$. All configurations will be defined on the whole $\mathbf{S}^1$ and are symmetric under the $\mathbb{Z}_2$ (working “upstairs” – in the terminology of⁷). This means, for example, that every time there is a 5-brane between the two fixed planes of the $\mathbb{Z}_2$ there is also a mirror 5-brane. There would be an equivalent formulation (“downstairs”) where configurations were only defined on the interval between the two “ends of the world”.

We know that M-theory on $\mathbb{R}^{9,1} \times \mathbf{S}^1/\mathbb{Z}_2$ is heterotic $E_8 \times E_8$ with one E_8 theory living on each fixed plane of the $\mathbb{Z}_2$. If we compactify this theory on K3 we need to supply a total of 24 instantons and 5-branes. The theory we are interested in is a single 5-brane coupled to an E_8 gauge theory. To achieve this we need to have no instantons in one of the E_8 theories and one 5-brane close to this “end of the world”. The remaining 23 instantons and 5-branes must therefore be either instantons in the other “end of the world” or 5-branes in the bulk.

In the 6-dimensional description the distance of the 5-brane from the “end of the world,” x , is a modulus. The effective gauge coupling of the E_8 depends on x . From the 6-dimensional point of view certain degrees of freedom connected to the 5-brane act as matter coupled to the E_8 gauge field. Since the couplings and masses of this matter depend on x , the low energy effective E_8 gauge coupling, g , will depend on x . Here we will calculate the x -dependence of g from M-theory or more precisely from 11-dimensional supergravity. For supergravity to be applicable all distances involved in the problem need to be much bigger than the 11-dimensional Planck scale. This means especially that $\text{Vol}(K3) \gg l_{Planck}^4$. Furthermore we are interested in the behaviour of the theory when it is close to the point with tensionless strings or equivalently with a zero size instanton, which is $x = 0$. To be in that situation we take $x \ll \text{vol}(K3)^{\frac{1}{4}}$. The x -dependence of the 6-dimensional gauge coupling g , comes about because the volume of the K3 at $X^{11} = 0$ depends on x .

To calculate g we need to find the form of the metric as a function of x . The calculation of the metric and of g is described in detail in¹. Here we will just state the result, which is

$$\frac{1}{g^2} = \frac{1}{g_0^2} - \frac{1}{8\pi^2} x T_2. \quad (6)$$

Here T_2 is the tension of the membrane in 11 dimensions. The expression xT_2 is the tension of the strings in the six-dimensional theory. This is because the membrane is stretched with one direction along the 11th direction and two directions along $\mathbb{R}^{5,1}$.

Compactifying further down to 4 dimensions on a torus of area A is straightforward

$$\frac{1}{g^2} = \frac{1}{g_0^2} - \frac{1}{8\pi^2} AxT_2. \quad (7)$$

This equation contains the needed information about the 4 dimensional theory. It tells how the gauge coupling in a E_8 gauge theory runs as a result of coupling to the 4 dimensional superconformal theory with E_8 global symmetry. In¹ it is shown how this implies that 10 of these saturate the β -function in complete agreement with the field theory derivation.

4 The 6D current-current correlator from F-theory

In this section, we use the duality between F theory on elliptic Calabi-Yau 3-folds and Heterotic String on $K3$ to compute the effective gauge coupling of heterotic string in six dimensions. We shall see that the result agrees completely with the corresponding M-theory calculation to first order. A second order effect which is suppressed by a factor of the volume of the $K3$ and by the length of $\mathbf{S}^1/\mathbb{Z}_2$ in calculations in the previous section naturally emerges in the F-theory setting. In the limit in which we extract the correlator for V_1 , i.e. taking the volume of $K3$ and the size of $\mathbf{S}^1/\mathbb{Z}_2$ to infinity, this second order effect vanishes.

We start with V_1 and couple it to a 6D E_8 SYM theory. The gauge theory is defined with a UV cut-off, but this imposes no problem for us since all we need is the dependence of the IR coupling constant on the VEVs of the V_1 theory. To be precise, we take the E_8 UV cut-off to be Λ and fix the E_8 coupling constant at Λ . The Coulomb branch of the V_1 theory has a single tensor multiplet. We denote the VEV of its scalar component by ϕ . ϕ is the tension of the BPS string in $\mathbb{R}^{5,1}$. In M-theory $\phi = xT_2$. The mass scale of the V_1 theory is thus $\phi^{1/2}$. We would like to find the dependence of the IR E_8 coupling constant on ϕ when $\phi \ll \Lambda$. Heuristically speaking, the running E_8 coupling constant will receive contributions from loops of modes from V_1 of mass $\sim \phi^{1/2}$.

The set-up that we have just described arises in the heterotic string compactified on $K3$ with a small E_8 instanton. We take the $(0, 23)$ embedding with a single 5-brane in the bulk close to the 9-brane with unbroken E_8 . The F-theory dual has a base B which is the Hirzebruch surface $\mathbf{F}_{11}$ with one point

blown-up^{10,6}. $\mathbf{F}_n$ is a $\mathbf{P}^1$ bundle over $\mathbf{P}^1$. Let the area of the fiber $\mathbf{P}^1$ in $\mathbf{F}_{11}$ (i.e. the Kähler class integrated over the fiber) be k_F and the area of zero section $\mathbf{P}^1$ of the fibration be k_D .

We blow-up a point in the zero section of the fibration of $\mathbf{P}^1$ over $\mathbf{P}^1$. There are 10 7-branes wrapping that zero-section and passing through a point of the exceptional divisor. These are responsible for the unbroken E_8 gauge group. Let k_E be the area of the exceptional divisor. The area of the above mentioned 7-brane locus (part with unbroken E_8) is k_D . The Kähler class is

$$k = (k_F - k_E)E + k_FD + (k_D + k_E - nk_F)F$$

where E, D, F are the cohomology classes of the exceptional divisor, base and fiber.

$$E \cdot E = -1, \quad E \cdot D = F \cdot D = 1, \quad (8)$$

$$D \cdot D = n - 1, \quad E \cdot F = F \cdot F = 0. \quad (9)$$

$$(10)$$

A 3-brane wrapping the exceptional divisor gives a BPS string in $\mathbb{R}^{5,1}$ (corresponding to the membrane connecting the M-theory 5-brane to the end of the world). Its tension is given by integrating the D3-brane tension over E . Using¹⁸

$$2\kappa^2\tau_p^2 = 2\pi(4\pi^2\alpha')^{3-p}$$

the tension of the BPS string is simply

$$\phi = \pi^{1/2}k_E$$

in the units $\kappa = 1$. The volume of the whole base is

$$V = \frac{1}{2}k \cdot k = k_F(k_D + k_E) - \frac{1}{2}k_E^2 - \frac{n}{2}k_F^2. \quad (11)$$

This volume is the 6D inverse gravitational constant and we have to keep it fixed. Although the V_1 modes have an effect on the gravitational constant as well, by dimensional analysis, this effect is much smaller than ϕ and behaves as $\sim \phi^2$. How should k_F depend on ϕ , in our setting? k_F measures the tension of 3-branes wrapped on F . On the heterotic side, these are elementary strings which occupy a point on K3. Their tension is fixed in the heterotic picture. Thus k_F is independent of ϕ .

Now we come to the gauge coupling. To do this calculation it is convenient to imagine that E_8 is broken down to $U(8) \subset E_8$. The gauge kinetic term for

8 unwrapped 7-branes of the same type is

$$\int \tau_7 \frac{(2\pi\alpha')^2}{4} \text{tr}_{\mathbf{8}}\{F^2\} d^8x.$$

We are working in the conventions

$$\text{tr}\{T^a T^b\} = \delta^{ab}, \quad a, b = 1 \dots 248.$$

For the $U(8)$ subgroup this means that

$$\text{tr}_{\mathbf{8}}\{T^a T^b\} = \frac{1}{2} \delta^{ab}.$$

This means that for a configuration of 10 7-branes forming an E_8 gauge theory the gauge kinetic term is:

$$\frac{1}{8} \int (2\pi\alpha')^2 \tau_7 \left(\sum_{a=1}^{248} F^a F^a \right) d^8x.$$

From this we read off (in units where $\kappa = 1$)

$$\frac{1}{4g^2} = \frac{1}{8} (2\pi\alpha')^2 \tau_7 = \frac{1}{32} \pi^{-3/2}.$$

Wrapping the 7-branes on D we get a 5+1D E_8 gauge theory with coupling constant

$$\frac{1}{4g^2} = \frac{1}{32} \pi^{-3/2} k_D.$$

From 11 we find that when V and k_F are kept fixed and $k_E = \pi^{-1/2} \phi$, the E_8 coupling constant is

$$\frac{1}{g(\phi)^2} = \frac{1}{8} \pi^{-3/2} [(k_D + k_E) - k_E] = \frac{1}{(g_0)^2} - \frac{1}{8\pi^2} \phi. \quad (12)$$

We have used the fact that $(k_D + k_E)$ is fixed to first order in ϕ when V is fixed. The other two terms in V are higher order corrections dual to taking $K3$ and the distance between the ends of the world to be large in the M-theory calculations. Eqn.12 describes the running of the E_8 coupling constant because of the coupling to V_1 . This is in complete agreement with the result 6 obtained from M-theory.

5 Discussion

We have found that for the 3+1D E_8 super-conformal theory with Seiberg-Witten curve

$$y^2 = x^3 + u^5,$$

the 2-point E_8 current correlator on the Coulomb branch satisfies:

$$\langle j_\mu^a(q) j_\nu^b(-q) \rangle = \begin{cases} \frac{C_2(E_8)}{40\pi^2} \delta^{ab} (q_\mu q_\nu - q^2 \eta_{\mu\nu}) \log\left(\frac{\Lambda}{|u|^{1/6}}\right) & \text{for } |q| \ll |u|^{1/6} \\ \frac{C_2(E_8)}{40\pi^2} \delta^{ab} (q_\mu q_\nu - q^2 \eta_{\mu\nu}) \log\left(\frac{\Lambda}{|q|}\right) & \text{for } |q| \gg |u|^{1/6} \end{cases}$$

where Λ is a UV cutoff which is an artifact of Fourier transforming.

We deduced that 10 copies of the E_8 theory can be coupled as “matter” to an $\mathcal{N} = 2$ E_8 SYM gauge field.

In 5+1D we found the expression for the low-energy limit of the 5+1D correlator of the $\mathcal{N} = (1, 0)$ E_8 theory on the Coulomb branch and away from the origin:

$$\langle j_\mu^a(q) j_\nu^b(-q) \rangle = -\frac{C_2(E_8)}{240\pi^2} \delta^{ab} (q^2 \eta_{\mu\nu} - q_\mu q_\nu) (\Lambda^2 - \phi) \quad \text{for } |q| \ll \phi.$$

where ϕ is the VEV of the scalar of the low-energy tensor multiplet.

It would be interesting to determine the correlator in the UV region $|q| \gg |\phi|$ or, equivalently, at the fixed point $\phi = 0$. It seems that the methods presented in this paper are not powerful enough for that purpose.

Acknowledgments

I wish to thank Yeuk-Kwan E. Cheung and Ori J. Ganor with whom this work was done. I furthermore wish to thank Steve Gubser, Igor Klebanov, Sangmin Lee, Sanjaye Ramgoolam and Savdeep Sethi for discussions. I wish to thank the organizers of the 1998 Trieste Conference on Superfivebranes and Physics in 5+1 Dimensions for letting me speak at the conference. This work was supported by the Danish Research Academy.

References

1. Yeuk-Kwan E. Cheung, Ori J. Ganor and Morten Krogh, “*Correlators of the global symmetry currents of 4d and 6D superconformal theories*,” *Nucl. Phys. B* **523**, 1988 (171), *hep-th/9710053*
2. E. Witten, “*Some Comments on String Dynamics*,” contributed to Strings ’95, *hep-th/9507121*

3. A. Strominger, "Open p -Branes," *Phys. Lett.B* **383**, 44 (1996), *hep-th/9512059*
4. E. Witten, "Strong Coupling Expansion Of Calabi-Yau Compactification," *Nucl. Phys. B* **471**, 135 (1996), *hep-th/9602070*
5. E. Witten, "Physical Interpretation Of Certain Strong Coupling Singularities," *Mod. Phys. Lett. A* **11**, 2649 (1996), *hep-th/9609159*
6. D.R. Morrison and C. Vafa, "Compactifications Of F-Theory On Calabi-Yau Threefolds - II," *Nucl. Phys. B* **476**, 437 (1996) *hep-th/9603161*
7. P. Horava and E. Witten, "Eleven-Dimensional Supergravity On a Manifold With Boundary," *Nucl. Phys. B* **475**, 94 (1996) *hep-th/9603142*
8. O.J. Ganor and A. Hanany, "Small E_8 Instantons and Tensionless Non-Critical Strings," *Nucl. Phys. B* **474**, 122 (1996), *hep-th/9602120*
9. O.J. Ganor, D.R. Morrison, N. Seiberg, "Branes, Calabi-Yau Spaces, and Toroidal Compactification of the $N = 1$ Six-Dimensional E_8 Theory," *Nucl. Phys. B* **487**, 93 (1997), *hep-th/9610251*
10. N. Seiberg and E. Witten, "Comments On String Dynamics In Six-Dimensions," *Nucl. Phys. B* **471**, 121 (1996), *hep-th/9603003*
11. C. Vafa, "Evidence For F-Theory," *Nucl. Phys. B* **469**, 403 (1996) *hep-th/9602022*
12. T. Banks, M.R. Douglas and N. Seiberg, "Probing F-theory With Branes," *Phys. Lett.B* **387**, 278 (1996) *hep-th/9605199*
13. N. Seiberg, "IR Dynamics on Branes and Space-Time Geometry," *Phys. Lett.B* **384**, 81 (1996), *hep-th/9606017*
14. J.D. Blum and K. Intriligator, "New Phases of String Theory and 6d RG Fixed Points via Branes at Orbifold Singularities," *Nucl. Phys. B* **506**, 199 (1997) *hep-th/9705044*
15. K. Intriligator, "New String Theories in Six Dimensions via Branes at Orbifold Singularities," *hep-th/9708117*
16. E. Witten, "Phase Transitions In M-Theory And F-Theory," *Nucl. Phys. B* **471**, 195 (1996) *hep-th/9603150*
17. N. Seiberg, "Five Dimensional SUSY Field Theories, Non-trivial Fixed Points AND String Dynamics," *Phys. Lett.B* **388**, 753 (1996) *hep-th/9608111*
18. J. Polchinski, "TASI Lectures on D-Branes," *hep-th/9611050*
19. S. Kachru and C. Vafa, "Exact Results For $N = 2$ Compactifications Of Heterotic Strings," *Nucl. Phys. B* **450**, 1995 (69), *hep-th/9505105*